

# START Modified ADULT

(size, + 2° sex characteristics)

Move the Walking Wounded **MINOR**

No Respirations after Head Tilt **EXPECTANT**

## CONTROL BLEEDING

Respiratory Distress (> 30/min) **IMMEDIATE**

Perfusion (No Radial Pulse) **IMMEDIATE**

Mental Status **IMMEDIATE**

(Unable to Follow Commands)

Normal RPM, Follows Commands **DELAYED**

## CONDUCT SECONDARY TRIAGE IN THE TREATMENT PHASE

### FL MCI LEVELS

MCI Level 1: 5-10 victims

MCI Level 2: 11-20 victims

MCI Level 3: 21-100 victims

MCI Level 4: 100 -1000 victims

MCI Level 5: Over 1000 victims

July 2021

# JumpSTART Modified

(Newborn to Young Adult\*)

Move the Walking Wounded **MINOR**

No Respirations and No Peripheral Pulse **EXPECTANT**

Respiratory Rate: > 45/min, < 15/min  
or †Work of Breathing, obvious distress **IMMEDIATE**

No Respirations with Peripheral Pulse  
Give 5 Ventilations via Barrier Device  
Spontaneous Respirations Resume  
after 5 Ventilations **IMMEDIATE**

No Spontaneous Respirations Resume  
after 5 Ventilations **EXPECTANT**

## CONTROL BLEEDING

Perfusion (No Palpable Pulse) **IMMEDIATE**

Mental Status\*\*  
Unresponsive or not localizing pain **IMMEDIATE**

Alert, responds to voice, localizes pain **DELAYED**

\*Presence of 2° sex characteristics; \*\*Consider developmental level  
July 2021 with permission ©Lou E Romig MD. [emlrc.org/flpedready/](http://emlrc.org/flpedready/)

CONDUCT SECONDARY TRIAGE IN THE TREATMENT PHASE